

Effects of *Scrophularia Striata* extracts on Wound Healing In Mice

Ali Akbar Jafari¹, Ali Mohamad Latify², Majid Shohrati³, Azad Aliahmadi⁴, Reza Hajhosseini⁵, Armin Hassanzadeh Inalou⁶, Amir Ali Shahbazfar⁷ and * Razzagh Mahmoudi⁸

¹ Department of Biology, Faculty of Science, Payam Noor University, Tehran, Iran,

² Applied Biotechnology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

³ Department of Clinical Pharmacy, Faculty of Pharmacy, Baqiyatallah University of Medical Sciences, Tehran, Iran,

⁴ Phytochemistry Research Center and Department of Pharmacognosy, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran,

⁵ Department of Biochemistry, Faculty of Science, Payam Noor University, Tehran, Iran,

⁶ DVM student, Faculty of Veterinary Medicine, University of Tabriz, Tabriz, Iran,

⁷ Associate professor, Department of Pathobiology, Faculty of Veterinary Medicine, University of Tabriz, Tabriz, Iran

⁸ Associate professor, Medical Microbiology Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

Abstract

Background and Aim: Study of different methods, to improve skin wounds healing caused by diabetes and burns has been always a matter of medical researches. Treatment with chemical drugs has its own problems so utilizing low-cost herbal treating methods has gained more attention day by day. The present research studied and compared the healing effects and functions of different *Scrophularia striata* extracts on mice wounds. Methods: At the first step the wound healing effect of three different kind of extracts (aqueous, alcoholic and hydroalcoholic) was investigated. Male mice weighting 25-35 gram were divided into five groups as follow: The First group considered as control (without treatment), the second group received phenytoin and the other three groups received 10 percent of aqueous, alcoholic and hydraulic extracts respectively. In order to control the wound healing process, wound's photos were documented in determined times and analyzed by image j software. In the second step, based on the first step findings, the wound healing effects of a hydroalcoholic extract ointment was investigated. In this step mice were divided into five groups including control, Ocerin treatment, phenytoin 1%, 5 percent and 10 percent herbal ointment. Results: Average times of wound healing were 9.1 ± 3.1 , 6.6 ± 1.9 , 6.5 ± 1.7 , 6.7 ± 1.8 , 6.1 ± 0.9 days orderly for the control group, Phenytoin, alcoholic, hydroalcoholic and aqueous extracts at the first step of the study. Hydroalcoholic extract had the best effect with the lowest healing time. At the second step healing times were 6 ± 0.6 , 6.43 ± 1.2 , 6.74 ± 2.04 , 8.86 ± 1.9 , 9.14 ± 2.09 days for 10 percent herbal ointment treatment, 5 percent herbal ointment treatment, Phenytoin treatment, ocerin treatment and control groups respectively. Conclusion: The study showed that 10% ointment had the most suitable effect compare with the other groups. This study showed that *Scrophularia Striata* can be one of the choices for manufacturing a medication for wound healing.

Keywords: Extract, mice, *scrophularia striata*, wound healing.

* Corresponding author: Medical Microbiology Research Center, Qazvin University of Medical Sciences, Qazvin, Iran